

BROWN ROOT ROT RESISTANCE

Test accepted: May 2008

Test updated: June 2024

Pathogen: *Phoma sclerotoides* G. Preuss ex Sacc.

Test authors: F.A. Gray, C.R. Hollingsworth, R.W. Groose, C.J. Reedy and R.C. Larsen

PLANT CULTURE

ContainerStuewe and Sons, Inc., Tall One Treepots™ (top) width = 10 cm, depth = 36 cm, vol. = 2.83 L (<https://www.stuewe.com>)

Plant Growth Medium.....Pasteurized soil (1 part sand/1 part sandy loam soil)

PlantingUse pregerminate seed and treat with *Sinorhizobium meliloti* prior to planting (<http://www.nitragin.com>)

No. of PlantsOne plant per pot and 24 plants per replicate

No. of Reps3 to 5 replications

Plant CareMaintain in the greenhouse at 21°C/15.6°C (day/night); 12 to 16 hr day length for 4 to 6 months. Since these pots are unstable, some type of support is required. We have used metal milk racks, which hold 12 pots each.

FUNGAL ISOLATION AND INOCULUM PRODUCTION

SourceBrown root rot (BRR) diseased alfalfa roots. Diseased roots in plastic bags can be stored under refrigeration (4°C) for 1 month.

Isolation*Phoma sclerotoides* can be isolated from diseased root tissue placed in water agar using standard isolation techniques. Plates must be incubated at 10°C for 2 months or until pycnidia are mature prior to sub-culturing.

ProductionCulture fungus in Petri plates containing potato dextrose agar maintained at 10°C until pycnidia are mature. Place 250 cc barley seed in a 600 ml glass beaker and add 130 ml water. Cover beaker with aluminum foil and autoclave for 60 min. Allow seed to cool overnight and reautoclave for an additional 60 min. When cool, transfer two mycelia plugs on to moistened seed and secure aluminum foil with parafilm. Maintain at 10°C for 2 months. Following thorough colonization of barley seed, remove and spread out on clean surface and allow to air dry.

StoragePlace colonized barley seed in plastic bags and store at -14°C. Inoculum should remain viable for up to 24 months or longer.

PLANT INOCULATION

Age of Plant4 to 6 months old

Type of Inoc.Infected barley seed containing mycelia and mature pycnidia

AmountTwo infected barley seeds/plant. Additional inoculum should increase disease development.

ProcedureInoculate plants in late summer to early fall. Remove soil from upper taproot. Place infected barley seeds against root, approximately 2.5 cm below the soil surface. Cover inoculum with a plug of sterile cotton and replace soil. Plant injury is not required.

INCUBATION OF INOCULATED PLANTS

CultureAfter inoculation, potted plants are placed outside on top of the ground and protected from cold temperatures by some type of insulation such as bales of hay to prevent roots from freezing. The experiment should have a rectangular design with only one bale in width to provide maximum protection. Loose hay should be placed into all openings to insure good insulation. Root infection has been reported to occur during host dormancy from early fall to early spring. Cardinal temperatures for fungal growth are 0°C (minimum), 15-16°C (optimum), and 27°C (maximum).

Plants should be watered as needed to maintain soil moisture. If straw bales do not provide sufficient insulation to prevent winterkill, one of the following two alternatives are suggested.

a) Place potted plants in the ground (dug-in) up to the top of the pots. A straw bale barrier of one bale high is still recommended for a windbreak, as well as the placement of loose straw over the plants, or

b) Field plots could be direct seeded, 6" row spacing, thinned to one plant every 6", inoculated as previously described, and evaluated the following spring. If this alternative is used, other plant pathogens and/or pests present in the test site may interfere with the results.

EvaluationPlants are left outside through the winter months. Plants can be evaluated for root rot in late spring to early summer. The fungus is inactive during the warm summer months. Since this test requires plants to be placed outside over one-half of the year, results may vary from year to year depending on local weather conditions.

RATING PLANTS FOR BROWN ROOT ROT

Evaluating plants for BRR severity. Partially remove plants from pots, remove soil from upper 10 cm of root, remove cotton plug and rinse under running water. Where potted plants have been 'dug-in' or where plants have been direct seeded, additional effort will be required for rating plants for BRR. Rate roots for disease response on a scale of 1-5 described below:

1No disease, roots healthy

2Slight disease, root tissues exhibit localized discoloration

3Moderate disease, well-defined lesions on root

4Severe root rot, plant deterioration

5Acute root rot, plant dead

CHECK CULTIVARS

Disease Reaction	Fall Dormancy Rating	Resistant Plants (%)*	
		Expected	Acceptable Range
Highly Resistant			
Peace (<i>Canada Dept. of Ag.</i>)	1	60	40-60
Moderately Resistant			
Multi-plier (<i>Mycogen Seeds</i>)	3	20	10-30

*Alfalfa populations can be characterized for reaction to brown root rot by the percent of resistant plants with scores of 1 and 2 compared to standard check cultivars.



(Click to see larger photo.)
Brown root rot symptoms.

CORRELATION TO FIELD TRIALS

A non-destructive forage yield trial, including the highly resistant and moderately resistant brown root rot check cultivars, was conducted at a site in Wyoming naturally infested with *P. sclerotioides*. Plant stands after three years were 86% for Peace and 16% for Multi-plier. An on-going repeat of the previously described trial conducted at a nearby *P. sclerotioides*-infested site, has produced similar but less dramatic results. Plant stands after three years for the BRR standard resistant and susceptible check cultivars were 97% for Peace and 67% for Multi-plier.

PATHOTYPES

Pathotypes of *Phoma sclerotioides* is available have not been reported. Studies by the authors have shown one Canadian (ATC #56515) and 13 Wyoming isolates of *P. sclerotioides* to all be pathogenic on alfalfa. Also, a *P. sclerotioides* isolate from cicer milkvetch (*Astragalus cicer* L.) was equally pathogenic on alfalfa when compared to an alfalfa isolate from Wyoming.

SOURCES OF INOCULUM

Deb Samac

USDA-ARS PSRU

1991 Upper Buford Circle

495 Borlaug Hall

St. Paul, MN 55108

(612) 625-1243

debby.samac@usda.gov

HELPFUL INFORMATION


Brown root rot of alfalfa is not a seedling disease. Good infection has occurred with four- to six-month-old plants. Also, disease development of inoculated plants has not been observed under normal greenhouse environmental conditions.

DISTRIBUTION

First reported on alfalfa in Canada in 1933, BRR of alfalfa has a circumpolar distribution in continental and alpine climates with severe winters. In North America, it occurs in Alaska, Yukon, the Northwest Territories, the four western provinces of Canada (British Columbia, Alberta, Saskatchewan, and Manitoba), as well as in Nova Scotia. First reported in the U.S. in Wyoming in 1969, it has since been reported from Idaho, Montana, Colorado and New Mexico in the West, Minnesota and Wisconsin in the Midwest, and New Hampshire, New York, and Vermont in the Northeast. To date, its distribution has occurred in Zones 1-4 of the USDA Plant Hardiness Zone Map.

<http://www.usna.usda.gov/Hardzone/ushzmap.html>

DISTRIBUTION AND SEVERITY OF BROWN ROOT ROT



-  States or provinces where BRR has been found on alfalfa but an extensive survey of alfalfa fields has not been made.
-  States or provinces where extensive surveys of alfalfa fields have been conducted or multiple reports have occurred.

(Click on the map above for a larger version.)

REFERENCES

1. Berkenkamp, B., S. Bittman, and D. McCartney. 1991. Resistance of alfalfa cultivars to brown root. *Can. J. Plant Sci.* 71:211-213.
2. Hollingsworth, C.R., Wunsch, M. J., and Bergstrom, G. C. 2015. Brown root rot. *Compendium of Alfalfa Diseases and Pests*, 3rd Edition. APS Press, St. Paul, MN. 138 pp.
3. Gray, F.A., C.R. Hollingsworth, C.J. Reedy, D.E. Legg, R.C. Larsen, R.W. Groose and D.W. Koch. 2008. Pathogenicity of fourteen isolates of *Phoma sclerotoides*, causing brown root rot of alfalfa. *Can. J. Plant Path.* Vol 30(2):285-293 (<https://doi.org/10.1080/07060661.2008.10540544>).
4. Gray, F.A., C.R. Hollingsworth, D.W. Koch, R.W. Groose, A.M. Gray, D.M. Woods, C.J. Reedy, T.E. Heald and J.R. Gill. 2003. Brown root rot of alfalfa. *Plant Science Timely information series*, No. 1, revised Feb. 2008. University of Wyoming, College of Agriculture, Department of Plant Sciences (<http://www.uwyo.edu/plants/publications/brrtimelyinfo2007update1.pdf>).
5. Gray, F.A., T.E. Heald, C.R. Hollingsworth and D.W. Koch. 1997. Brown root rot caused by *Phoma sclerotoides*, a new disease of alfalfa in the U.S. p. 22-24. In *Proc. 10th West Alfalfa Improvement Conf.*, Davis, CA, 27-28 June 1997.
6. Hollingsworth, C.R. and F.A. Gray. 1999. First report of brown root rot on alfalfa caused by *Phoma sclerotoides* in the continental United States. *Plant Disease* 83:1071 (Disease Notes).
7. Hollingsworth, C.R. 1999. Biology and management of brown root rot, *Phoma sclerotoides* of alfalfa. M.Sc. thesis, University of Wyoming, Laramie, WY, 82071, pp. 1-110.
8. Hollingsworth, C.R. 2002. Assessing heritability of brown root rot (*Phoma sclerotoides*) resistance and forage yield in nine alfalfa (*Medicago sativa* ssp. *sativa*) populations. Ph.D. dissertation, University of Wyoming, Laramie, WY 82071, pp. 1-133.
9. Hollingsworth, C.R., F.A. Gray and T.E. Heald. 2000. Pathogenicity of *Phoma sclerotoides*, a new disease in the continental U.S. *Proceedings/ Reports of the American Forage and Grassland Council and the 37th North American Alfalfa Improvement Conference*, p. 299. Madison, WI. July 16-19, 2000.
10. Hollingsworth, C.R., F.A. Gray, D.W. Koch, R.W. Groose and T.E. Heald. 2003. Distribution of *Phoma sclerotoides* and incidence of brown root rot in Wyoming, U.S.A. *Can. J. of Plant Path.* 25:215-217.
11. Hollingsworth, C.R., F.A. Gray and R.W. Groose. 2005. Evidence for the heritability of resistance to brown root rot of alfalfa caused by *Phoma sclerotoides*. *Can. J. Plant Path.* 27:64-70.
12. Hollingsworth, C.R., R.W. Groose and F.A. Gray. 2003. Comparative disease reactions of Cycle 0 and Cycle 1 alfalfa following inoculation with *Phoma sclerotoides*. *Phytopathology* 93:536 (Abstract).