

MULTIFOLIOLATE LEAF EXPRESSION (LEAVES WITH GREATER THAN 3 LEAFLETS/LEAF)

March 1995

C.C. Sheaffer, M. McCaslin, J.J. Volenec, J.H. Cherney, K.D. Johnson, W.T. Woodward, and D.R. Viands

FIELD ESTABLISHMENT

SeedingPlants should be established in rows using transplants (8-12 week old) or by direct seeding and thinning. Procedures for greenhouse culture of plants for transplanting are provided in the procedures for fall dormancy determination (see fall dormancy testing procedures, Reference 1).

Spacing30 cm between plants in rows 0.6 to 1.0 m apart.

No. of Plants25 per replication

No. of Reps4 minimum

OtherMaintain a high soil fertility, a soil pH of 6.5 to 7.5, and adequate soil moisture. Minimize weed and insect interference. Inoculate with *Rhizobium meliloti*.

RATING

Clip herbage using locally accepted schedules and rate plants were vegetative in late summer. Multifoliolate leaf (MF) expression on a per plant basis is visually scored. The multifoliolate leaf expression score per plant is based on an average of individual stems and on number of MF leaves per stem:

Score	Description
0	= all trifoliolate leaves
1	= 1 MF leaf/stem
2	= 2-3 MF leaves/stem
3	= 4-5 MF leaves/stem
4	= 6-7 MF leaves/stem
5	= 8+ MF leaves/stem

CALCULATIONS

MF Expression IndexPlants should be established in rows using transplants (8-12 week old) or by direct seeding and thinning. Procedures for greenhouse culture of plants for transplanting are provided in the procedures for fall dormancy determination (see fall dormancy testing procedures, Reference 1).

$$\text{MFI} = \frac{\Sigma[(\text{No. } 1 \times 1) + (\text{No. } 2 \times 2) + (\text{No. } 3 \times 3) + (\text{No. } 4 \times 4) + (\text{No. } 5 \times 5)]}{\text{Total No. plants in population}}$$

Both the MFI value and the percentage of plants with MF expression (those plants scoring 1 to 5) should be provided when describing experimental germplasm and cultivars.

CHECK CULTIVARS

Category	Cultivar	MFI	Acceptable Range
Trifoliolate	Vernal	1.00	1.00-1.05
Low MF	Legend	1.86	1.40-2.40
Moderate MF	MultiKing I	2.55	2.00-3.00
High MF	Proof	3.35	2.80-3.80

SCIENTISTS WITH EXPERTISE

Craig C. Sheaffer

University of Minnesota
Dept of Agronomy and Plant Genetics
411 Borlaug Hall
St. Paul, MN 55108
(612) 625-7225

J. J. Volenec

Purdue University
Department of Agronomy
Lilly Hall of Life Sciences
West Lafayette, IN 47907
(317) 494-8071

REFERENCES

1. Barnes, D.K., D.M. Srnith, L.R. Tueber, and M.A. Peterson. 1991. Fall dormancy. In C.C. Fox et al., (ed) Standard tests to characterize alfalfa cultivars. North American Alfalfa Improvement conf.
2. Juan, N.A., C.C. Sheaffer, and D.K. Barnes. 1993. Temperature and photoperiod effects on multifoliolate expression and morphology of alfalfa. *Crop Sci.* 33:573-578.
3. Etzel, M.G., J.J. Volenec, and J.J. Vorst. 1988. Leaf morphology, shoot growth, and gas exchange of multifoliolate alfalfa phenotypes. *Crop Sci.* 28:263-269.
4. Volenec, J.J., and J.H. Cherney. 1990. Yield components, morphology, and forage quality of multi-foliolate alfalfa phenotypes. *Crop Sci.* 30:1234-1238.