

Developing Practical Phosphorus & Potassium Tissue Test Recommendations & Utilizing Struvite in Modern Alfalfa Systems III

*O. Steve Norberg, Washington State University
Steve C. Fransen, Washington State University
Joseph H. Harrison, Washington State University
Donald A. Llewellyn, Washington State University
Elizabeth Whitefield, Washington State University
Jon Paul Driver, Washington State University*

Developing critical plant nutrient levels in-season improves recommendations and applications, saving producers time, expense and effort since many growers take samples for hay quality. Three experiments were designed as follows: 1) Phosphorus (P) Rate study with differing rates of P_2O_5 using monoammonium phosphate (MAP); including: 0, 30, 60, 120, 240 lb P_2O_5 acre⁻¹ on a low testing P soil <10 ppm (Olsen P method); 2) Potassium (K) Rate study with differing rates of K_2O using potassium sulfate: 0, 40, 80, 160, 240, 320 lb K_2O acre⁻¹ on an <100 ppm K soil (ammonium acetate method). The following is summation of four years of results for alfalfa harvested at mid-bud stage for all cuttings in the same field. Increasing P rate from 0 to 240 lb P_2O_5 acre⁻¹ increased yield by 0.9, 1.5, 1.6 and 1.5 tons acre⁻¹ in 2018, 2019, 2020 and 2021, respectively. Averaged over 2019 & 2020 and assuming \$560 ton⁻¹ for 11-52-0 P_2O_5 (\$0.538 lb P_2O_5 ⁻¹), the whole plant tissue level at the economic optimum was 0.35 and 0.36% at mid-bud stage for 150 and \$225 ton⁻¹ of hay, respectively. About 80% of yield response occurred in the first and second cuttings indicating that P and K need to be applied in the fall or early spring to get the largest yield response. Potassium concentration in alfalfa varied widely between years optimum K content for the two years but the optimum for \$200 ton⁻¹ hay, when the price lb of K_2O is was \$0.36, is 1.9 and 1.6% for 2019 and 2020, respectively. The relationship between potassium alfalfa tissue concentration and yield was not significant. Both the price of fertilizer and the price of hay both impact the recommended rate of P and K significantly and must be taken into account when determining optimum fertilizer rates in the current economic market and we have developed an excel spreadsheet to help with these recommendations.